



Leucidal[®] SF MAX

Patent Pending: Application Number 14/740,887

Code Number: FSSM15019MAX

INCI Name: *Lactobacillus* Ferment

INCI Status: Approved

REACH Status: Fully Compliant

CAS Number: 1686112-36-6

EINECS Number: N/A

Origin: Biotechnology | *Lactobacillus*

Processing:

GMO Free

No Ethoxylation | No Irradiation | No Sulphonation

No Ethylene Oxide treatment | No Hydrogenation

Additives: None | **Preservatives:** None | **Antioxidants:** None

Other additives: None

Solvents used: Water

Appearance: Clear to Hazy, Colorless to Yellow Liquid

Soluble/Miscible: Water Soluble

Suggested Use Levels: 2.0 - 4.0%

Suggested Applications: Skin Conditioning, Antimicrobial



BACKGROUND

You have spoken, and they listened. Leucidal[®] SF MAX is the latest iteration of our Leucidal[®] line. Optimizing the fermentation process to tighten the specification for bacteriocins, Leucidal[®] SF MAX provides broad spectrum antimicrobial results with an increased degree of confidence.

Consumer choice is the most important factor when it comes to cosmetic sales. Today, a growing number of consumers are opting to move away from synthetic preservatives such as parabens, formaldehyde donors and phenoxyethanol. In addition to public pressure, the use of these synthetic materials in cosmetics is also becoming more strictly regulated. For these reasons, formulators have been actively searching for alternatives to synthetic preservatives that can provide broad spectrum antimicrobial activity and can be added into a wide range of cosmetic applications.

A full line of products were developed and derived from naturally occurring compounds that provide broad spectrum antimicrobial protection. As a result, these novel natural antimicrobials are considered self-preserving cosmetic actives and therefore can be used as consumer-friendly alternatives to synthetic preservatives in a wide range of cosmetic applications.

SCIENCE

Leucidal[®] SF MAX is a powerful probiotic-based ingredient created by the fermentation of *Lactobacillus* in a defined growth medium. *Lactobacillus* is one of the species of microorganisms used to produce fermented products, such as kimchi and sauerkraut, a Korean dietary staple, from cabbage. Like many members of the lactic acid bacteria family, *Lactobacillus* is capable of restricting the growth of other microorganisms by acidifying its environment. However, *Lactobacillus* also produces novel antimicrobial peptides, known as bacteriocins, that are capable of providing broad spectrum antimicrobial protection. During the manufacturing process, lysozyme is added to the ferment filtrate to facilitate a controlled cell lysis to ensure the release of the antimicrobial peptides for maximized activity.

Information contained in this technical literature is believed to be accurate and is offered in good faith for the benefit of the customer. The company, however, cannot assume any liability or risk involved in the use of its chemical products since the conditions of use are beyond our control. Statements concerning the possible use of our products are not intended as recommendations to use our products in the infringement of any patent. We make no warranty of any kind, expressed or implied, other than that the material conforms to the applicable standard specification. Freedom from patent infringement is not implied. All information is for investigative purposes only.



BENEFITS

A skin moisturization study was performed using an untreated control, generic cream base, and an experimental with the same cream base containing 2.0% Leucidal® SF Max. Comparative moisturization results from this study are shown in Figure 1.

When compared to the base cream Leucidal® SF Max improved moisturization by 14.38% and after 24 hours and by 25.00% after four weeks. Based on these results, adding this innovative product provides the formulator the opportunity to capitalize on both the natural antimicrobial properties of Leucidal® SF Max, as well as its ability to provide potent moisturizing benefits to the cosmetic formulation.

Comparative Moisturization




Figure 1. Percent Difference in Moisturization for **Leucidal® SF Max**

Leucidal® SF Max was designed to maximize the effectiveness of the antimicrobial peptides in our original Leucidal® Liquid SF. The ability of Leucidal® SF Max to inhibit the growth of a variety of bacteria and fungi was determined using the Minimum Inhibitory Concentration (MIC) test. The results are illustrated in Figure 2, showing that this material provides broad spectrum antimicrobial protection.

Microorganism Tested	MIC (%)
<i>E. coli</i>	0.50
<i>P. aeruginosa</i>	0.50
<i>S. aureus</i>	0.50
<i>A. brasiliensis</i>	0.50
<i>C. albicans</i>	0.50

Figure 2. MIC data for **Leucidal® SF Max**.

The positive MIC screening results warranted further testing to confirm its ability to provide product preservation. Double Challenge Tests were completed using either 2.0% or 4.0% Leucidal® SF Max in a generic cream base formulation at pH values of 3, 5, and 7. Samples were inoculated with *E. coli*, *P. aeruginosa*, *S. aureus*, *C. albicans*, and *A. brasiliensis*.

Information contained in this technical literature is believed to be accurate and is offered in good faith for the benefit of the customer. The company, however, cannot assume any liability or risk involved in the use of its chemical products since the conditions of use are beyond our control. Statements concerning the possible use of our products are not intended as recommendations to use our products in the infringement of any patent. We make no warranty of any kind, expressed or implied, other than that the material conforms to the applicable standard specification. Freedom from patent infringement is not implied. All information is for investigative purposes only.



During the first 28-day incubation period, samples were periodically collected and tested for the presence of these microorganisms. Following this initial 28 days of incubation, the cream samples were then re-inoculated with the microbial cultures and sampled over an additional 28-day period. Figure 3 shows the positive preservation results for Leucidal® SF Max tested at 4.0% in a generic cream base formulation at pH 5.

4.0% Leucidal® SF Max in Cream Formula Challenge Test - pH 5




Figure 3. Challenge Test results for Generic Cream Formula pH 5 with 4.0% Leucidal® SF Max inoculated on Day 0 and re-inoculated on Day 28. Results show log reduction in viable organisms.

A Time Kill Test was performed to determine the change in population of aerobic microorganisms within a specified sampling time when tested against 4.0% Leucidal® SF Max solution. The activity of the test material inoculated was evaluated at determine time intervals of 30 seconds, 1, 5, 10 and 30 minutes after the inoculation to determine quantitatively the number of viable microorganisms remaining after the incubation time. As shown in Figure 4, the Gram-positive and Gram-negative bacteria as well as the yeast and mold were reduced by 99.9% within 30 seconds interval of the test after the inoculation.

Information contained in this technical literature is believed to be accurate and is offered in good faith for the benefit of the customer. The company, however, cannot assume any liability or risk involved in the use of its chemical products since the conditions of use are beyond our control. Statements concerning the possible use of our products are not intended as recommendations to use our products in the infringement of any patent. We make no warranty of any kind, expressed or implied, other than that the material conforms to the applicable standard specification. Freedom from patent infringement is not implied. All information is for investigative purposes only.



4.0% Leucidal® SF Max Time Kill Test




Figure 4. Time Kill Test results for 4.0% Leucidal® SF Max.

USE RECOMMENDATIONS

Leucidal® SF Max can be used in a wide range of cosmetic products, however to ensure optimum results we recommend using the following guidelines. Incorporate the product into formulations at a pH between 3 and 8, during the cooling phase of the process at temperatures lower than 70°C.

Information contained in this technical literature is believed to be accurate and is offered in good faith for the benefit of the customer. The company, however, cannot assume any liability or risk involved in the use of its chemical products since the conditions of use are beyond our control. Statements concerning the possible use of our products are not intended as recommendations to use our products in the infringement of any patent. We make no warranty of any kind, expressed or implied, other than that the material conforms to the applicable standard specification. Freedom from patent infringement is not implied. All information is for investigative purposes only.